

Low-energy Effective Theory for One-dimensional Lattice Bosons near Integer Filling

Yu-Wen Lee,¹ Yu-Li Lee,² and Min-Fong Yang¹

¹*Department of Physics, Tunghai University, Taichung, Taiwan*

²*Department of Physics, National Changhua University of Education, Changhua, Taiwan*

(Dated: February 6, 2008)

A low-energy effective theory for interacting bosons on a one-dimensional lattice at and near integer fillings is proposed. It is found that two sets of bosonic phase fields are necessary in order to explain the complete phase diagram. Using the present effective theory, the nature of the quantum phase transitions among various phases can be identified. Moreover, the general condition for the appearance of the recently proposed Pfaffian-like state can be realized from our effective action.

PACS numbers: 05.30.Jp, 03.75.Kk, 03.75.Lm, 71.10.Pm, 75.10.Jm

I. INTRODUCTION

Much effort has been devoted to understanding the effects of competing interactions on quasi-one-dimensional systems. Recent advances in loading ultracold bosonic atoms into an optical lattice lead to the realization of one-dimensional (1D) lattice boson systems, and inspire the investigation on many-body quantum phenomena therein.^{1,2,3,4} Besides the nearest-neighbor hopping t and the tunable on-site interaction U , a sizable nearest-neighbor interaction V is now within experimental reach by using the dipolar interaction among atoms.^{5,6} These experiments raise the interest in creating and detecting exotic quantum phases in 1D interacting lattice boson systems.

For 1D lattice bosons at integer filling, the phase diagram obtained by the mean-field calculation includes three different phases. They are the Mott insulator (MI) for large U , a charge density wave (CDW) for large V , and a superfluid (SF) for large t .^{7,8} However, by using the Density Matrix Renormalization Group method, it is found recently that, at filling of $\bar{n} = 1$, there exists a novel insulating phase lying between the CDW and MI phases in the weak coupling region.⁹ The phase transition between this new phase and the conventional insulating phases (i.e., the CDW and MI phases) is found to be of second order. Similar to the Haldane phase of quantum spin-one chains,¹⁰ this new insulating phase can be identified by a highly nonlocal string order parameter, and thus called as the Haldane insulator (HI) phase.⁹ Another investigation employing quantum Monte Carlo simulations shows that, when $t \ll U \lesssim 2V$, there exists a supersolid (SS) phase lying between the $\bar{n} = 1$ CDW and the $\bar{n} = 1/2$ CDW phases.¹¹ The CDW-SS transition is found to be of second order with dynamic critical exponent $z = 2$. Recently, due to the possibility of a fault tolerant quantum computation based on non-abelian anyons,¹² people begin to search for models containing non-abelian anyons,¹³ as well as for techniques for their detection and manipulation.¹⁴ It is pointed out that bosonic atoms in a 1D lattice with infinite repulsive three-body on-site interactions around $\bar{n} = 1$ have a ground state very close to a gapless Pfaffian-like state,¹⁵

which may serve as the basis to create non-abelian anyons in 1D systems.

With an eye on these new developments, it is desirable to have a unified understanding to the rich phases and the quantum phase transitions among them through a suitable low-energy effective theory. A common way of deriving low-energy effective theory for 1D bosonic systems is to use Haldane's density-phase representation (also known as "phenomenological bosonization").^{16,17} Such an approach has been successfully used to address the low-energy physics of SF and the related Mott transitions at commensurate fillings.¹⁷ However, this effective theory contains *only one* set of collective variables which describe long-wavelength phase and density fluctuations, and can show only one phase transition between two phases. That is, it fails to show two phase transitions found in Ref. 9. Moreover, in this simplest one-component hydrodynamic effective action, the CDW and the SF order parameters compete (or, are dual to) each other. Therefore, from such an effective theory, it is hard to understand the origin of the SS phase reported in Ref. 11, where the CDW and the SF orders coexist. In the present work, a low-energy effective theory with *two* sets of bosonic phase fields is proposed. Within the present framework, not only the nature of the phase transitions and the corresponding critical modes can be explicitly identified, the phase diagrams reported in Refs. 9 and 11 can also be easily understood. Moreover, we find that the SS phase in 1D has at most an *algebraic* long-range CDW order and has an algebraic *boson-paired supersolid* order instead of the ordinary one. Furthermore, the general condition for the appearance of the gapless Pfaffian-like state can be realized within the present approach. This observation will be of help for searching gapless Pfaffian-like state in more realistic systems. We believe that the effective theory shown below provides a general description of the low-energy physics for 1D interacting lattice bosons near integer filling, and it serves as a starting point for further investigations on the static and dynamical properties of these systems.

II. THE EXTENDED BOSON HUBBARD MODEL AND ITS EFFECTIVE THEORY

The lattice bosons with a nearest-neighbor hopping t , an on-site interaction U , and a nearest-neighbor interaction V can be described by the extended Bose-Hubbard model (EBHM)

$$H = -t \sum_{\langle i,j \rangle} (b_i^\dagger b_j + h.c.) + \frac{U}{2} \sum_i \delta n_i^2 + V \sum_{\langle i,j \rangle} \delta n_i \delta n_j, \quad (1)$$

where $b_i^\dagger$ is the bosonic creation operator at site i , $n_i = b_i^\dagger b_i$ is the number operator, and $\delta n_i \equiv n_i - \bar{n}$ measures deviations of the particle number from a mean filling $\bar{n}$. Since our main concern is in discussing the phase transition among various insulating phases, where the local particle number fluctuations should not be strong, near an integer filling n_0 (i.e., $\bar{n} \simeq n_0$), one can truncate the bosonic Fock space to only three local states with particle numbers n_0 and $n_0 \pm 1$.^{9,15,18,19,20} Now the lattice boson model in Eq. (1) can be mapped onto a spin-one chain in the truncated Hilbert space,

$$\begin{aligned} H_{\text{spin}} = & -\frac{n_0 t}{2} \sum_{\langle i,j \rangle} (S_i^+ S_j^- + h.c.) \\ & + V \sum_{\langle i,j \rangle} S_i^z S_j^z + \frac{U}{2} \sum_i (S_i^z)^2 \\ & - \frac{n_0 t \xi}{2} \sum_{\langle i,j \rangle} [S_i^z S_i^+ S_j^- + S_i^- S_i^z S_j^+ + S_i^- S_j^z S_j^+ \\ & + S_i^+ S_j^- S_j^z + \xi (S_i^z S_i^+ S_j^- S_j^z + S_i^- S_i^z S_j^z S_j^+)] \end{aligned} \quad (2)$$

due to the correspondence

$$b_i^\dagger \leftrightarrow \sqrt{\frac{n_0}{2}} (1 + \xi S_i^z) S_i^+, \quad \delta n_i \leftrightarrow S_i^z. \quad (3)$$

S_i^α are the $S = 1$ spin operators, and the parameter $\xi \equiv \sqrt{(n_0 + 1)/n_0} - 1$ is a measure of the ‘particle-hole asymmetry’. For simplicity, ξ is set to zero, which is valid at least when $n_0 \gg 1$. From the viewpoint of the bosonization theory, the main physics is indeed not modified by the effect due to the ξ -dependent terms.²¹ By further using the composite spin representation for the $S = 1$ spin operators,^{22,23} where they are rewritten as a sum of two commuting spin-1/2 species, the problem of spin-one chain becomes that with two coupled spin-1/2 chains. After applying the Jordan-Wigner transformation and bosonization procedure for these spin-1/2 operators, one finally arrives at the following bosonized effective

Hamiltonian^{22,23,24}

$$\begin{aligned} H_{\text{eff}} = & \frac{u_s}{2} \int dx \left[K_s (\partial_x \Theta_s)^2 + \frac{1}{K_s} (\partial_x \Phi_s)^2 \right] \\ & + g_1 \int dx \cos(\sqrt{8\pi} \Phi_s) \\ & + \frac{u_a}{2} \int dx \left[K_a (\partial_x \Theta_a)^2 + \frac{1}{K_a} (\partial_x \Phi_a)^2 \right] \\ & + \int dx \left[g_2 \cos(\sqrt{8\pi} \Phi_a) + g_3 \cos(\sqrt{2\pi} \Theta_a) \right], \end{aligned} \quad (4)$$

where Φ_s , Θ_s and Φ_a , Θ_a are the symmetric (called as the charge mode) and the antisymmetric (called as the neutral mode) combinations of the bosonized fields for the two spin-1/2 operators, respectively. The bosonized forms of the original lattice boson operators are

$$\begin{aligned} \frac{b_j^\dagger}{\sqrt{a}} & \sim \sqrt{\frac{n_0}{2}} \sqrt{\frac{2}{\pi a}} e^{i\sqrt{\frac{\pi}{2}} \Theta_s} \cos\left(\sqrt{\frac{\pi}{2}} \Theta_a\right) + \dots, \\ \frac{\delta n_j}{a} & \sim \sqrt{\frac{2}{\pi}} \partial_x \Phi_s - (-1)^{x/a} \frac{2}{\pi a} \sin \sqrt{2\pi} \Phi_s \cos \sqrt{2\pi} \Phi_a \end{aligned} \quad (5)$$

with $x = ja$, where a is the lattice constant and plays the role of a short-distance cutoff. In the derivation of Eq. (4), some irrelevant (or less relevant) terms have been dropped. As a result, the charge and the neutral modes become decoupled. We note that, while the method of phenomenological bosonization can give identical form of the charge part in Eq. (4),¹⁷ the neutral mode is missed in that approach. That is the reason why the conventional theory fails to account for the complete phase diagram of the interacting lattice bosons near the integer fillings. As shown in Refs. 22 and 23, assuming the couplings in the two coupled spin-1/2 chains being weak, the values of the parameters in Eq. (4) can be obtained as follows: $g_1 = -g_2 = (2V - U)/2\pi^2 a$, $g_3 = -n_0 t / \pi a$; the Luttinger parameters $K_s = [1 + (U + 6V)/\pi n_0 t]^{-1/2}$, $K_a = [1 - (U - 2V)/\pi n_0 t]^{-1/2}$; and the sound mode velocities $u_s = u_0/K_s$, $u_a = u_0/K_a$ with $u_0 \equiv n_0 t a$. Since the present effective Hamiltonian is reached under several approximations, the values of the above parameters should not be treated very seriously. Here we would like to consider Eq. (4) as a phenomenological theory. It provides a universal low-energy physics of the 1D interacting lattice boson models near integer fillings, such that the nature of the phase transitions and the long-distance behaviors of the major correlation functions at each phase can be predicted. As shown below, this effective Hamiltonian does explain the results reported recently.^{9,11,15} It reinforces our belief in the validity of the present framework.




FIG. 1: The schematic phase diagram at integer filling $\bar{n} = n_0$ in the weak-coupling regime. The phase boundaries of Kosterlitz-Thouless, Gaussian, and Ising types are denoted by the solid, dashed, and dotted lines, respectively.

III. THE PHASES AND PHASE TRANSITIONS OF THE EXTENDED BOSON HUBBARD MODEL

Through the discussion on the relevance of each non-linear term in Eq. (4), the phase diagram can be obtained as shown in Fig. 1, which looks similar to that reported in Ref. 9 in the weak-coupling regime. According to the work in Ref. 22, better agreement can be reached if one keeps more local states (say, with particle numbers n_0 , $n_0 \pm 1$, and $n_0 \pm 2$) and maps the interacting lattice boson model to a spin chain with higher integer spin.²⁶ However, we must emphasize that one of the most important feature of the bosonization analysis in Ref. 22 is that the properties of the corresponding spin model do not dramatically depend on the value of the spin quantum number S , but only on whether S is integer or half-odd integer. This suggests that the main results obtained below from our analysis of the Hamiltonian in Eq. (4) should remain unchanged even when the local particle number fluctuations become stronger. The phase boundary separating the SF and the MI phases is determined by $K_s = 1$. This comes from the relevance of the g_1 term in the charge sector of Eq. (4), and thus gives a type of Kosterlitz-Thouless transition.²⁷ The MI-HI phase transition is given by $g_1 = 0$. It is a Gaussian transition, which separates two gapped phases. The HI-CDW transition occurs when $K_a = 1/2$. It is due to the competition of the two relevant terms (i.e., the g_2 and the g_3 terms) in the neutral sector of Eq. (4), and thus is a Ising-type transition.²⁸ By using the bosonized forms of the original lattice boson operators in Eq. (5), the correlation functions at long distance can be calculated. In the SF phase, because the charge mode is gapless while the neutral field Θ_a is pinned, the density fluctuation is gapless and the single-particle correlation function shows an algebraic long range order (LRO) with an enhanced superfluid correlation: $\langle b_i^\dagger b_j \rangle \sim |i - j|^{-1/4K_s}$. True LRO can develop when $K_s \rightarrow \infty$. In the CDW phase, both of Φ_s and Φ_a are pinned. In this case, while

	SF	MI	HI	CDW
K_s	> 1	< 1	< 1	< 1
K_a	$> \frac{1}{2}$	$> \frac{1}{2}$	$> \frac{1}{2}$	$< \frac{1}{2}$
g_1	irrelevant	< 0	> 0	> 0
g_2	irrelevant	irrelevant	irrelevant	< 0
$\langle \Phi_s \rangle$	not pinned	0	$\sqrt{\pi/8}$	$\sqrt{\pi/8}$
$\langle \Phi_a \rangle$	not pinned	not pinned	not pinned	0
$\langle \Theta_a \rangle$	0	0	0	not pinned

TABLE I: The Luttinger parameters and the roles of various couplings in different phases at integer fillings. The values of the fields $\Phi_{s,a}$ and Θ_a indicated in this table are the values at which these fields were pinned in each phase. Notice that throughout our discussions, we take the coupling $g_3 < 0$ (following Ref. 22), and it is irrelevant only in the CDW phase above.




FIG. 2: The schematic phase diagram near integer filling $\bar{n} \simeq n_0$. The solid lines denote the commensurate-incommensurate transition with $z = 2$.

the single-particle correlation functions decay exponentially, the staggered part of the density-density correlation shows LRO: $\langle \delta n_i \delta n_j \rangle \sim (-1)^{|i-j|}$. This justifies the identification of the CDW phase. In the MI and the HI phases, both Φ_s and Θ_a are pinned, and both the single-particle and density correlation functions decay exponentially at long distance. Both in these two gapped phases, the lattice translation symmetry does not break. The only difference is that the Φ_s field is pinned at 0 and $\sqrt{\pi/8}$ in the MI and the HI phases, respectively. As pointed out in Ref. 9, one can distinct the HI phase from the MI one via a nonlocal string order parameter defined by $\mathcal{O} \sim \langle \delta n_i \exp(i\pi \sum_{k=i}^j \delta n_k) \delta n_j \rangle$. Within the present bosonization approach, such a string order parameter can be expressed as $\mathcal{O} \sim \langle \sin(\sqrt{\pi/2}\Phi_s(x_i)) \sin(\sqrt{\pi/2}\Phi_s(x_j)) \rangle$,²⁹ which is completely independent of the neutral mode. Thus it is indeed nonzero in the HI phase with Φ_s being pinned at $\langle \Phi_s \rangle = \sqrt{\pi/8}$, consistent with the findings in Ref. 9.³⁰ To make the relations between the relevant couplings and the pinned fields in each phase more explicit, we summarize the above discussions in table I.

Up to now, our discussion has focused on systems exactly at integer fillings. The deviation from integer fill-

ing can be studied by turning on a nonzero chemical potential $\delta\mu$ and adding a term $\delta\mu \sum_i \delta n_i$ in the Hamiltonian in Eq. (1). ($\delta\mu$ is measured from the middle of the lobe.) In the bosonized representation, this term takes the form $\delta\mu \sqrt{\frac{2}{\pi}} \partial_x \Phi_s$, which affects the charge mode only.³¹ The $\partial_x \Phi_s$ term tends to create a finite number of kinks in the Φ_s configuration, and it will compete with the Sine-Gordon term $\cos \sqrt{8\pi} \Phi_s$ which tends to pin down Φ_s . When $\delta\mu$ is comparable to the magnitude of the charge gap of the insulating phases, such a term will trigger a well-known commensurate-incommensurate transition.¹⁷ As long as K_s is not too small such that the breather mass gap is larger than the Sine-Gordon kink mass gap, the universal features of this transition is well captured by the Luther-Emery limit of the charge mode, where the bosonized effective Hamiltonian can be further mapped to a free massive Dirac fermion theory with a chemical potential term.²⁵ At the transition point $\delta\mu = \delta\mu_c$ such that the charge gap closes, the gapless charge mode excitation density obeys the scaling relation $\rho \sim (\delta\mu - \delta\mu_c)^{1/2}$, and the dynamic critical exponent of this transition becomes $z = 2$. After the transition, the charge mode is described by a Gaussian model with nonzero stiffness, which accounts for the SF behavior in 1D. Therefore, starting from either the MI or the HI phase, nonzero $\delta\mu$ can induce a transition to the SF phase. On the other hand, after the transition from the CDW phase, even though the gapless charge mode shows the SF behavior, the CDW-like behavior can remain, and thus the SS phase is obtained. It is because the neutral sector is unaffected and the Φ_a field is still pinned in this case even after the transition. According to this analysis, it is tempted to interpret the SS phase near integer fillings as the descendant of the CDW phase at integer fillings. Thus our effective theory does explain the occurrence of a $z = 2$ second-order CDW-SS phase transition found in Ref. 11. It is noted that, since the charge mode is gapless in the SS phase, a power-law behavior in the density-density correlation function $(-1)^{|i-j|} \langle \delta n_i \delta n_j \rangle \propto |i-j|^{-K_s}$ is expected. That is, the SS phase has at most an *algebraic* long-range CDW order. Because $K_s \ll 1$ in the SS phase, such a power-law decay may not be examined easily by finite-size numerical calculations. Another interesting feature of the SS phase is that, while the superfluid stiffness is nonzero, the single-particle correlation $\langle b_i^\dagger b_j \rangle$ decays *exponentially* due to pinning of the Φ_a field (or the Θ_a field being disorder). However, the boson-pair correlation $\langle (b_i^\dagger)^2 b_j^2 \rangle$ indeed shows an algebraic long range order. Thus the system behaves somewhat like a “boson-paired superfluid”. This issue deserves further examinations by other approaches. The above discussions are summarized by the phase diagram shown in Fig. 2.

IV. THE PFAFFIAN-LIKE STATES

Besides the application to the 1D EBHM, a recently proposed gapless Pfaffian-like state¹⁵ as the ground state of three-body-hard-core bosons in 1D lattice can also be understood within the present framework. Due to the infinite three-body on-site repulsion, only three local states with occupation numbers $n = 0, 1, 2$ are necessary. Similar to the previous discussion, the interacting lattice boson model in Ref. 15 can again be mapped to a spin-one chain in Eq. (2) with $U = V = 0$ and $n_0 = 1$. That is, the lattice boson model studied in Ref. 15 is just a special case of the lattice spin-one Hamiltonian studied in the present paper. Therefore, the proposed gapless Pfaffian-like state should correspond to one of the *gapless* ground states described by the present continuum effective field theory. When $U = V = 0$ and $n_0 = 1$, ignoring the ξ -dependent terms in Eq. (2) as we did before, the low-energy effective Hamiltonian becomes that in Eq. (4) with the Luttinger parameters $K_s = 1$ and $K_a > 1/2$. Thus it indicates that the gapless Pfaffian-like state should be identified as the ground state laying on the SF-MI phase boundary, where the charge sector is gapless and the Θ_a field is pinned. As a nontrivial cross-check of this observation, we show that the correlation functions calculated by the Pfaffian-like wave function under a completely different approach can indeed be reproduced within the present framework. By using the bosonized forms Eq. (5) of the original lattice boson operators, for the ground state laying on the SF-MI phase boundary, the single-particle and boson-pair correlation functions behave at long distance

$$\langle b_i^\dagger b_j \rangle \sim |i-j|^{-1/4K_s}, \quad \langle (b_i^\dagger)^2 b_j^2 \rangle \sim |i-j|^{-K_s} \quad (6)$$

with $K_s = 1$. Notice that, on the SF-MI phase boundary, the exponent of the single-particle correlation function is always *four times larger* than that of the boson-pair correlation function. These results agree with those in Ref. 15. It supports our observation that the ground state laying on the SF-MI phase boundary is nothing but the Pfaffian-like state. Since a direct strong three-body interaction among atoms are rare in nature,³² the authors in Ref. 15 suggested to use a Feshbach-resonant system away from the resonance to generate an effective three-body interaction among atoms, thus realizing this novel state. According to the present discussion, we find that strong three-body on-site repulsion is not essential for the appearance of the Pfaffian-like state. Instead, 1D Interacting lattice boson models can have a gapless Pfaffian-like ground state, as long as they are tuned toward the critical points of the SF-MI phase transition. Thus it is plausible to realize such 1D Pfaffian-like states in an ordinary boson Hubbard model without borrowing the mechanism of Feshbach resonance.

V. CONCLUSIONS

To summarize, a low-energy description, which successfully capture the rich phases of the 1D lattice boson model near integer fillings, is constructed. We believe that the present effective theory provides general applications to related problems. For example, parallel approaches can be used to study models of coupled 1D hard-core lattice bosons, where similar effective theories can be constructed.^{33,34,35,36,37,38} When local density fluctuations become stronger, more local states with different particle numbers should be kept. Following the

above reasoning, an effective theory with more bosonic phase fields should be reached. It is interesting to see if more novel phases can be found in this case.

Acknowledgments

Y.-W. Lee, Y.-L. Lee, and M.-F. Yang acknowledge the support by the National Science Council of Taiwan under the contracts NSC 96-2112-M-029-006-MY3, NSC 96-2112-M-018-006-MY2, and NSC 95-2112-M-029-006, respectively.

-
- ¹ B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch, *Nature* (London) **429**, 277 (2004).
 - ² T. Kinoshita, T. Wenger, and D. S. Weiss, *Science* **305**, 1125 (2004).
 - ³ T. Stöferle, H. Moritz, C. Schori, M. Köhl, and T. Esslinger, *Phys. Rev. Lett.* **92**, 130403 (2004).
 - ⁴ C. D. Fertig, K. M. O'Hara, J. H. Huckans, S. L. Rolston, W. D. Phillips, and J. V. Porto, *Phys. Rev. Lett.* **94**, 120403 (2005).
 - ⁵ A. Griesmaier, J. Werner, S. Hensler, J. Stuhler, and T. Pfau, *Phys. Rev. Lett.* **94**, 160401 (2005); A. Griesmaier, J. Stuhler, T. Koch, M. Fattori, T. Pfau, and S. Giovanazzi, *ibid.* **97**, 250402 (2006).
 - ⁶ *Special issue on Ultracold Polar Molecules: Formation and Collisions*, *Eur. Phys. J. D.* **31** (2004).
 - ⁷ K. Goral, L. Santos, and M. Lewenstein, *Phys. Rev. Lett.* **88**, 170406 (2002).
 - ⁸ D. L. Kovrizhin, G. V. Pai, and S. Sinha, *Europhys. Lett.* **72**, 162 (2005).
 - ⁹ E. G. Dalla Torre, E. Berg, and E. Altman, *Phys. Rev. Lett.* **97**, 260401 (2006).
 - ¹⁰ F. D. M. Haldane, *Phys. Lett.* **93A**, 464 (1983); *Phys. Rev. Lett.* **50**, 1153 (1983).
 - ¹¹ G. G. Batrouni, F. Hébert, and R. T. Scalettar, *Phys. Rev. Lett.* **97**, 087209 (2006).
 - ¹² A. Kitaev, *Ann. Phys. (N.Y.)* **303**, 2 (2003); M. H. Freedman, M. J. Larsen, and Z. Wang, *Commun. Math. Phys.* **227**, 605 (2002).
 - ¹³ A. Kitaev, *Ann. Phys. (N.Y.)* **321**, 2 (2006).
 - ¹⁴ S. Das Sarma, M. H. Freedman, and C. Nayak, *Phys. Rev. Lett.* **94**, 166802 (2005); A. Stern and B. I. Halperin, *ibid.* **96**, 016802 (2006); P. Bonderson, A. Kitaev, and K. Shtengel, *ibid.* **96**, 016803 (2006).
 - ¹⁵ B. Paredes, T. Keilmann, and J. I. Cirac, *Phys. Rev. A* **75**, 053661 (2007).
 - ¹⁶ F. D. M. Haldane, *Phys. Rev. Lett.* **47**, 1840 (1981).
 - ¹⁷ T. Giamarchi, *Quantum Physics in One Dimension* (Oxford University Press, New York, 2004).
 - ¹⁸ E. Altman and A. Auerbach, *Phys. Rev. Lett.* **89**, 250404 (2002).
 - ¹⁹ S. D. Huber, E. Altman, H. P. Büchler, and G. Blatter, *Phys. Rev. B* **75**, 085106 (2007).
 - ²⁰ As shown in Refs. 18 and 19, the derived theories under such a truncation of the Hilbert space can even successfully describe the SF-MI transition. Thus the restriction of allowing local particle number fluctuation of order one should be sufficient to capture the correct critical theory in the whole phase diagram.
 - ²¹ Their irrelevance can be realized as follows. First, the forms of the leading nonlinear terms in our effective action Eq. (4) are severely constrained by the global U(1) symmetry, the lattice translation symmetry and the compactification radius of the bosonic phase fields. Therefore, the ξ -dependent terms should not affect them. Second, while bosonized terms like $\partial_x \Phi_s \partial_x \Theta_a$ or $\partial_x \Phi_s$ are indeed generated by the ξ -dependent terms, these bosonized terms will either modify the Luttinger parameters (since the neutral mode is gapped) or their effects are already included in the chemical potential term. Hence, we conclude that the ξ -dependent terms will not make any qualitative change for present discussions as long as $\xi \ll 1$.
 - ²² H. J. Schulz, *Phys. Rev. B* **34**, 6372 (1986).
 - ²³ J. Sólyom and J. Timonen, *Phys. Rev. B* **34**, 487 (1986).
 - ²⁴ We follow here the conventions used in Ref. 25, where free fermions are mapped to bosonic phase fields with a compactification radius $R = 1/\sqrt{4\pi}$.
 - ²⁵ A. M. Tsvelik, *Quantum Field Theory in Condensed Matter Physics*, 2nd ed. (Cambridge University Press, Cambridge, 2003).
 - ²⁶ In fact, our Fig. 1 is nothing but Fig. 1 in Ref. 22 after translating to the language of lattice boson model. When more local states are kept, the phase diagram will correspond to Fig. 2 in Ref. 22. We notice that at a qualitative level, Fig. 2 and Fig. 1 of Ref. 22 are almost identical, i.e. the spin one case and the other integer spin cases possess the same phases apart from the slight changes of phase boundaries.
 - ²⁷ J. M. Kosterlitz and D. J. Thouless, *J. Phys. C* **6**, 1181 (1973); J. M. Kosterlitz, *ibid.* **7**, 1046 (1974).
 - ²⁸ This can be most easily seen through a “refermionization” procedure where we introduce a set of right- and left-moving Dirac fermions $\chi_{R,L} = \frac{1}{\sqrt{2\pi a}} e^{\pm i\sqrt{4\pi}\phi'_{R,L}}$ with $\phi'_{R,L} = \frac{1}{\sqrt{2K_a}} \Phi_a \mp \sqrt{\frac{K_a}{2}} \Theta_a = \Phi_a \mp \sqrt{\frac{1}{4}} \Theta_a$ to rewrite the Hamiltonian. For further details of the bosonization description of the Ising transition, we refer the reader to the book in Ref. 25.
 - ²⁹ The derivation of this form has some subtle aspects, which are mainly attributable to the continuous-field (coarsegrained) scheme. M. Nakamura [*Physica B* **329-333**, 1000 (2003)] has resolved such an ambiguity of the

bosonized expressions using a symmetry cleverly.

- ³⁰ Since Φ_s is also pinned at $\langle \Phi_s \rangle = \sqrt{\pi}/8$ in the CDW phase, the string order parameter is also nonvanishing there, consistent with the findings shown in Ref. 9.
- ³¹ We notice that although the staggered part of the chemical potential term (see Eq. (5)) does not enter into the continuum effective theory at the tree level, at the second order of perturbation theory, it can generate a term proportional $-(\delta\mu)^2 \cos(\sqrt{8\pi}\Phi_a)$ which renormalizes the coupling g_2 towards *more negative values*. This will certainly enhance the density wave ordering tendency, and hence, suggests that the SS phase analyzed below for small $\delta\mu$ should maintain its robustness in a finite region away from integer fillings.
- ³² Realizations of the three-body contact interactions and exotic quantum Hall states (including the bosonic nonabelian Pfaffian state) in a *three-dimensional* rapidly rotating Bose condensate are discussed in N. R. Cooper, Phys. Rev. Lett. **92**, 220405 (2004).
- ³³ P. Donohue and T. Giamarchi, Phys. Rev. B **63**, 180508(R) (2001).
- ³⁴ T. Stoferle, H. Moritz, C. Schori, M. Kohl, and T. Esslinger, Phys. Rev. Lett. **92**, 130403 (2004).
- ³⁵ M. A. Cazalilla, A. F. Ho, and T. Giamarchi, New. J. of Phys. **8**, 158 (2006).
- ³⁶ L. Mathey, A. Polkovnikov, and A. H. Castro Neto, arXiv:cond-mat/0612174.
- ³⁷ Yu-Wen Lee and Yu-Li Lee, Phys. Rev. A **73**, 043606 (2006).
- ³⁸ Yu-Wen Lee, unpublished.